

ANATOMICAL DIAGRAMS

FOR THE USE OF ART STUDENTS.

*ARRANGED WITH ANALYTICAL NOTES
AND DRAWN OUT BY*

JAMES M. DUNLOP, A.R.C.A.,

ANTIQUE AND LIFE CLASS MASTER AND LECTURER ON ARTISTIC
ANATOMY IN THE GLASGOW SCHOOL OF ART.

WITH INTRODUCTORY PREFACE BY

JOHN CLELAND, M.D., LL.D., F.R.S.,

PROFESSOR OF ANATOMY IN THE UNIVERSITY OF GLASGOW.

FOURTH EDITION.

New York
THE MACMILLAN COMPANY

1919

All rights reserved

FA 6478.5


HARVARD COLLEGE LIBRARY
GIFT OF
BERNARD BERENSON
Oct 13, 1938

To

**THE CHAIRMAN AND GOVERNORS OF THE
GLASGOW SCHOOL OF ART,
IN WHOSE SCHOOL I HAVE BEEN LECTURER
ON ARTISTIC ANATOMY
FOR A NUMBER OF YEARS ;
TO THE HEADMASTER MR. FRANCIS H. NEWBERY
AND TO THE STUDENTS,
THIS BOOK IS RESPECTFULLY DEDICATED
BY THEIR OBEDIENT SERVANT,**

JAMES M. DUNLOP.

GLASGOW, 1899.



P R E F A C E.

SCIENCE AND ART are indeed sisters, but they are very different in their tastes, and it is no easy task to cultivate with advantage the favour of both. Artistic Anatomy is in its nature a scientific pursuit, dealing partly in explicit observation of details of form, partly in the explanation of the causes producing them; while the details themselves are among those with which the followers of Art require to be familiar; and are sometimes of little apparent scientific importance save from an artistic point of view. In these circumstances it is little to be wondered at that this department of knowledge has not been more fully explored.

Properly conceived of, Artistic Anatomy undertakes the systematic study of the particulars of superficial form, the accurate description of them one by one, and the investigation of the structural and functional causes on which they depend.

Among the phenomena to be considered, the proportions of the great divisions of the body one to another claim an important place, and have justly received attention from remote times. Rules have been laid down by which an ideal standard has been sought to be fixed, the deviations produced by age and sex being taken into account; and while such standards are more or less artificial, and not to be too slavishly followed to the extent of an unnatural uniformity, they certainly are invaluable as expressing a mean which cannot be deviated from to more than a limited extent without transgressing the laws of nature and producing deformity.

Each part of the body has also its particular proportions, and the study of proportions passes gradually into that of details of shape. All these details are capable of being taken one by one and systematically described. But this cannot be done either accurately or instructively without reference to the subjacent structures on which they depend, and the actions governing the conditions of such structures.

Subcutaneous prominences of bone afford so many constant points in the surface of the figure, while the softer subcutaneous tissues sometimes occur in masses of such firmness as to be but little affected by change of attitude, and in other instances are flaccid, pendulous, wrinkled or stretched. But the muscles and their tendons produce the greatest variations of local form in different persons and in different attitudes; muscular substance swelling when in action, while tendons are incapable alike of swelling and of altering their total length, but may stand out when they are tightened over the concavity formed by the bending of a joint. Also lines of attachment to subcutaneous bone, themselves incapable of change of form, may in different circumstances be prominent or sunk according to the degree of swelling of the muscles around. Besides all this it must be noted that muscular contractions cause, especially in the face, lines, elevations, and depressions, not corresponding to the shapes of the muscles, but produced by the displacement of skin and subcutaneous fat, as illustrated by the elevation of the cheek and lower eyelid in laughter, and by the formation at the same time of the lines called crows' toes, and it does seem possible that a more careful analysis than has been attempted of the lines and displacements occurring in different expressions might yield better results than are to be obtained from such works as those of Le Brun, Sir Charles Bell, Piderit and Darwin. however valuable these may be. It may also be mentioned that

considerations in connection with balance, respiration, mental capacity and race fall within the scope of Artistic Anatomy.

If these views are allowed to be correct, it will be admitted that the field of Artistic Anatomy has never been covered; and if this task be ever undertaken it must be for its own sake, aiming at independent completeness, and not at mere assistance to Artists. Much will thus be brought to light, in all probability now unsuspected, and Art and Philosophy will both be gainers.

While, however, Art is one thing and Artistic Anatomy quite another, and while it is to be acknowledged that beautiful representations may be achieved without any anatomical knowledge, this only shows how much can be done by practised observation led on by intuitive appreciation which, often unconsciously, guides the mind to the accomplishment of its aims. But such success is neither easy nor to be depended on, and the general average thus obtainable cannot be expected to be so good as would be obtained if observation were assisted by acquaintance with the meanings of the shapes observed. The greatest masters, including notably Michelangelo, Leonardo da Vinci and Raphael, have found that to give intelligence to their efforts at representation, and enable them to understand the indispensable relations of parts it was necessary to call in the aid of dissection. For the eye, though often, even when well trained, at fault, especially when invention is brought into play, is yet subtle to detect instinctively the unsatisfactoriness of error.

It seems sometimes to be supposed that Artistic Anatomy is merely Anatomy made easy for Artists by omitting explicit details and all mention of internal organs,—superficial Anatomy in both senses of the word. But what is superficial in the sense of being slovenly is of little use to any one. The Professional Anatomist addressing his discourse to Artists, and desiring to give them the information for which they crave, cannot help

seeing at once that there is much internal structure which can have no possible bearing on Art, but he will fail altogether in his purpose if he does not note that the artist seeks for direction with regard to details which are often of small interest to the surgeon, and have received little attention from Anatomists.

Two of the sets of considerations most important to the Artist will easily be seen to be, one, the part played by the skeleton in determining the external form, and another, the precise extent and attachments of superficial muscles, together with the disposition of muscular fibre and tendon in individual muscles. It is principally to these two considerations that Mr. Dunlop directs attention in the following pages, appealing to the eye, instead of depending on description; and it appears to me that the method which he has selected, and the manner in which he has carried it out, provide for the Art Student a singularly compendious and desirable book, easily consulted, and occupying ground which has not hitherto been taken up. It is not the whole subject of Artistic Anatomy, but only one department of it which is here dealt with. The facts taught are brought out with diagrammatic simplicity and precision which cannot fail to bring them clearly and prominently before the student, thus giving him immense assistance. I have pleasure therefore in anticipating for this useful work a great success.

JOHN CLELAND.

CONTENTS

	PAGE
SIDE VIEW OF THE HEAD AND NECK—Bones	2
SIDE VIEW OF THE HEAD AND NECK—Muscles	3
FRONT VIEW OF THE HEAD AND NECK—Bones	4
FRONT VIEW OF THE HEAD AND NECK—Muscles	5
SIDE VIEW OF THE TRUNK—Bones	6
SIDE VIEW OF THE TRUNK—Muscles	7
FRONT VIEW OF THE TRUNK—Bones	8
FRONT VIEW OF THE TRUNK—Muscles	9
BACK VIEW OF THE TRUNK—Bones	10
BACK VIEW OF THE TRUNK—Muscles	11
OUTER VIEW OF THE LOWER LIMB—Bones	12
OUTER VIEW OF THE LOWER LIMB—Muscles	13
FRONT VIEW OF THE LOWER LIMB—Bones	14
FRONT VIEW OF THE LOWER LIMB—Muscles	15
INNER VIEW OF THE LOWER LIMB—Bones	16
INNER VIEW OF THE LOWER LIMB—Muscles	17
BACK VIEW OF THE LOWER LIMB—Bones	18
BACK VIEW OF THE LOWER LIMB—Muscles	19
OUTER VIEW OF THE UPPER LIMB—Bones	20
OUTER VIEW OF THE UPPER LIMB—Muscles	21
FRONT VIEW OF THE UPPER LIMB—Bones	22
FRONT VIEW OF THE UPPER LIMB—Muscles	23
INNER VIEW OF THE UPPER LIMB—Bones	24
INNER VIEW OF THE UPPER LIMB—Muscles	25
BACK VIEW OF THE UPPER LIMB—Bones	26
BACK VIEW OF THE UPPER LIMB—Muscles	27
THE BONES IN RELATION TO THE OUTLINE OF FIGURE—Front View	28
CONSTRUCTION LINES OF THE STANDING FIGURE—Front View	29
THE BONES IN RELATION TO THE OUTLINE OF FIGURE—Back View	30
CONSTRUCTION LINES OF THE STANDING FIGURE—Back View	31
THE BONES IN RELATION TO THE OUTLINE OF FIGURE—Side View	32
PARTS OF THE BONES WHICH DIRECTLY AFFECT THE SURFACE FORM—Side View	33
THE BONES AS IN ACTION—Front View	34
THE MUSCLES AS IN ACTION—Front View	35
THE BONES AS IN ACTION—Back View	36
THE MUSCLES AS IN ACTION—Back View	37
THE BONES AS IN ACTION—Side View	38


CONTENTS

	PAGE
THE MUSCLES AS IN ACTION—Side View	39
DETAILS OF THE FACE	40
MUSCLES OF THE HEAD, with notes of their attachments	41
MUSCLES OF THE NECK, with notes of their attachments	42
MUSCLES OF THE TRUNK, with notes of their attachments	43
MUSCLES OF THE TRUNK, with notes of their attachments	44
MUSCLES OF THE TRUNK, with notes of their attachments	45
MUSCLES OF THE TRUNK, with notes of their attachments	46
MUSCLES OF THE TRUNK, with notes of their attachments	47
MUSCLES OF THE TRUNK, with notes of their attachments	48
MUSCLES OF THE TRUNK, with notes of their attachments	49
MUSCLES OF THE THIGH—Front View, with notes of their attachments	50
MUSCLES OF THE THIGH—Front View, with notes of their attachments	51
MUSCLES OF THE BUTTOCK AND HIP, with notes of their attachments	52
MUSCLES OF THE THIGH—Back View, with notes of their attachments	53
MUSCLES OF THE LEG—Back View, with notes of their attachments	54
MUSCLES OF THE LEG—Front View, with notes of their attachments	55
OUTER VIEW OF THE FOOT—Bones and Muscles, with notes of their attachments	56
INNER VIEW OF THE FOOT—Bones and Muscles, with notes of their attachments	57
MUSCLES OF THE ARM—Front View, with notes of their attachments	58
MUSCLES OF THE ARM—Back View, with notes of their attachments	59
MUSCLES OF FRONT AND INNER SIDE OF FOREARM, with notes of their attachments	60
MUSCLES OF THE BACK AND OUTER SIDE OF FOREARM, with notes of their attachments	61
BONES OF THE HAND—Palmar View, and Muscles of Palm	62
MUSCLES OF THE HAND—Palmar View, and attachments	63
BONES OF THE HAND—Back View	64
MUSCLES OF THE HAND—Back View	65
SURFACE FORMS OF THE LIMBS	66
BONES OF THE UPPER AND LOWER LIMBS—Photos	67
DIAGRAMMATIC NOTES OF VARIOUS BONES	68
SURFACE FORMS OF THE TRUNK	69
THREE-QUARTER FRONT VIEW OF SKELETON—Photo	70
THREE-QUARTER BACK VIEW OF SKELETON—Photo	71
DIAGRAMS WITH NOTES ON PROPORTION	72

ANATOMICAL DIAGRAMS.


**SIDE VIEW OF THE
HEAD AND NECK.**

BONES.




SIDE VIEW OF THE HEAD AND NECK.

MUSCLES.




FRONT VIEW OF THE HEAD AND NECK.

BONES.




**FRONT VIEW OF THE
HEAD AND NECK.**

MUSCLES.




SIDE VIEW OF
TRUNK.

BONES.




**SIDE VIEW OF
THE TRUNK.**

MUSCLES.




FRONT VIEW OF THE TRUNK

BONES




FRONT VIEW OF
THE TRUNK.

MUSCLES.




**BACK VIEW OF
THE TRUNK.**

BONES.




BACK VIEW OF
THE TRUNK.

MUSCLES.




OUTER VIEW OF
THE LOWER LIMB.

BONES.




OUTER VIEW OF
THE LOWER LIMB.

MUSCLES.




FRONT VIEW OF
THE LOWER LIMB.

BONES.




**FRONT VIEW OF
THE LOWER LIMB**

MUSCLES.




INNER VIEW OF THE LOWER LIMB

BONES




INNER VIEW OF
THE LOWER LIMB.

MUSCLES.




**BACK VIEW OF
THE LOWER LIMB.**

BONES.




BACK VIEW OF
THE LOWER LIMB.

MUSCLES.




**OUTER VIEW OF
THE UPPER LIMB.**

BONES.




**OUTER VIEW OF
THE UPPER LIMB.**

MUSCLES.




**FRONT VIEW OF
THE UPPER LIMB.**

BONES.




FRONT VIEW OF
THE UPPER LIMB.

MUSCLES.




**INNER VIEW OF
THE UPPER LIMB.**

BONES.




INNER VIEW OF
THE UPPER LIMB.

MUSCLES.




**BACK VIEW OF
THE UPPER LIMB.**

BONES.




BACK VIEW OF THE UPPER LIMB

MUSCLES.



THE BONES IN RELATION TO THE OUTLINE OF FIGURE FRONT VIEW



CONSTRUCTION LINES OF THE STANDING FIGURE. FRONT VIEW




Fig. 1




Fig. 2

In the above diagrams, Fig. 1 shows the leading constructive lines when the figure is standing upright and resting on both legs. In this position the line passing through the shoulders, and that drawn across the antr. supr. spines of the Pelvis, are both obviously at right angles to the vertical axis of the body.


In Figure 2 the weight of the body is carried mainly on one leg, and in this position the axial line of the body becomes a curve, but still the line drawn through the shoulders and that through the Pelvis may be regarded as at right angles to this imaginary curve. Observe that the Pelvis is higher on the side which supports the figure, and also that the hip makes a sharper angle on that side, and farther, note the slope of the standing leg with the ground, necessary for the balance of the figure.

The axial line of the body carried down through the standing leg gives here a line of double curvature, which is the first line to be drawn in suggesting the pose.

THE BONES IN RELATION TO OUTLINE OF FIGURE.
BACK VIEW.




CONSTRUCTION LINES OF THE STANDING FIGURE. BACK VIEW.




The above diagrams show the leading constructive lines of the back view of the figure in similar positions to those already shown in the front view. The greater length of the trunk as viewed from behind will be observed on comparison with the front view. The lines across the back are drawn through the shoulders, the lower end of the thorax, the iliac crests, and the gluteal fold.

THE BONES IN RELATION TO OUTLINE OF FIGURE.
BACK VIEW.




CONSTRUCTION LINES OF THE STANDING FIGURE. BACK VIEW.




The above diagrams show the leading constructive lines of the back view of the figure in similar positions to those already shown in the front view. The greater length of the trunk as viewed from behind will be observed on comparison with the front view. The lines across the back are drawn through the shoulders, the lower end of the thorax, the iliac crests, and the gluteal fold.

THE BONES IN RELATION TO OUTLINE OF FIGURE.
SIDE VIEW.



**PARTS OF THE BONES WHICH DIRECTLY AFFECT
THE SURFACE FORM—SIDE VIEW.**



References to the bones

HEAD

1. Bones of the cranium; the shaded part is more or less plainly revealed upon the surface
2. Nasal bone
3. Malar or cheek-bone and Zygomatic arch
4. Mastoid process of Temporal bone
5. Lower jaw-bone, outline of its entire length

TRUNK


6. Clavicle
7. Spine of Scapula
8. Base of Scapula
9. Sternum
10. Cartilages of Ribs
11. Iliac Crest
12. Anterior superior iliac spine
13. Pubis

LOWER LIMB

14. Great trochanter of Femur
15. Patella
16. Outer condyle of Femur
17. Head of Tibia
18. Head of Fibula
19. Outer malleolus of Fibula
20. Os Calcis
21. Tuberosity of 5th metatarsal
22. Metatarsal bones

NOTE.—The parts of the Bones and Cartilages which are subcutaneous, or sufficiently near the surface to affect the surface form, are in this diagram marked in blue.

THE BONES AS IN ACTION—FRONT VIEW.



References to the Bones and Cartilages in the three views of the figure in action.

TRUNK.

- A—Clavicle
- B—Acromion process of Scapula
- B'—Spine of Scapula
- B''—Base of Scapula
- C—Sternum
- D—Cartilages of the Ribs
- E—Anterior superior iliac spine
- F—Iliac crest
- G—Posterior superior iliac spine
- H—Pubis
- I—Sacrum

LOWER LIMB.

- J—Great trochanter of Femur
- K—Outer condyle of Femur
- L—Inner condyle of Femur
- M—Patella
- N—Head of Tibia
- N'—Shaft of Tibia
- O—Head of Fibula
- P—Outer malleolus of Fibula
- Q—Inner malleolus of Tibia
- R—Os Calcis
- S—Tuberosity of 5th metatarsal
- T—Ball of great toe

UPPER LIMB.

- U—Inner condyle of Humerus
- V—Outer condyle of Humerus
- W—Olecranon process of Ulna
- X—Posterior border of Ulna
- Y—Head and styloid process of Ulna
- Z—Styloid process of Radius

NOTE.—The parts of the Bones and Cartilages which are subcutaneous or sufficiently near the surface to affect the surface form directly, are, in this diagram marked in blue. The parts so marked are therefore of great importance in sketching out the masses of the figure.

THE MUSCLES AS IN ACTION—FRONT VIEW.



References to the Muscles
in the three views of the
figure in action

HEAD AND TRUNK

- | | |
|--------------------------------------|---------------------|
| 1. Sternomastoid | 7. Rectus abdominis |
| 2. Posterior triangle
of the neck | 8. Serratus magnus |
| 3. Trapezius | 9. Latissimus dorsi |
| 4. Pectoralis major | 10. Infra-spinatus |
| 5. Deltoid | 11. Teres minor |
| 6. External oblique | 12. Teres major |
| | 13. Rhomboid |

LOWER LIMB

- | |
|-------------------------------|
| 14. Gluteus medius |
| 15. Gluteus maximus |
| 16. Tensor fasciæ femoris |
| 17. Sartorius |
| 18. Rectus femoris |
| 19. Vastus externus |
| 20. Vastus internus |
| 21. Iliacus and Psoas |
| 22. Pectineus |
| 23. Adductor longus |
| 24. Gracilis |
| 25. Adductor magntus |
| 26. Biceps femoris |
| 27. Semitendinosus |
| 28. Semimembranosus |
| 29. Plantaris |
| 30. Gastrocnemius |
| 31. Soleus |
| 32. Peroneus longus |
| 33. Peroneus brevis |
| 34. Peroneus tertius |
| 35. Extensor longus digitorum |
| 36. Tibialis anticus |
| 37. Extensor brevis digitorum |
| 38. Abductor minimi digiti |
| 39. Abductor pollicis |

UPPER LIMB.

- | |
|-------------------------------------|
| 40. Biceps |
| 41. Triceps |
| 42. Brachialis anticus |
| 43. Coraco-brachialis |
| 44. Supinator longus |
| 45. Extensor carpi radialis longior |
| 46. " " brevior |
| 47. " communis digitorum |
| 48. " ossis metacarpi pollicis |
| 49. " primi internodii pollicis |
| 50. " secundi internodii pollicis |
| 51. " carpi ulnaris |
| 52. Anconeus |
| 53. Flexor carpi ulnaris |
| 54. Flexor sublimis digitorum |
| 55. Palmaris longus |
| 56. Flexor carpi radialis |
| 57. Pronator teres |



FASCIÆ, APONEUROSES, &c.

- | |
|---------------------------------|
| a. Aponeurosis covering Rectus |
| b. Lumbar aponeurosis |
| c. Fascia lata—ilio-tibial band |
| d. Quadriceps |
| e. Ligamentum patellæ |
| f. Tendo-Achilles |

DETAILS OF THE FACE.



CARTILAGES OF THE NOSE.



MUSCLES OF THE HEAD.




Diagram showing **TEMPORALIS** muscle from temporal ridge and fossa above, and attached below to the coronoid process of the lower jaw-bone, the Zygomatic arch being supposed removed





Diagram showing **MASSETER** muscle attached to the Zygomatic arch and to the lower jaw-bone



II. Hyoid bone
 T. Thyroid cartilage of the Larynx
 (Pomum Adami)
 T.G. Thyroid gland
 Tr. Trachea or windpipe


References to muscles of the face

1. **ORBICULARIS PALPEBRARUM**
 from the tendo-palpebrarum, the frontal bone and superior maxillary, at the inner margin of orbit; it blends with occipito-frontalis and other muscles
2. **CORRUGATOR SUPERCILII**
 from frontal bone at the internal angular process; it blends with occipito-frontalis
3. **PYRAMIDALIS NASI**
 a small slip prolonged downwards from the occipito-frontalis to the nasal bones
4. **COMPRESSOR NARIS**
 from superior maxillary bone to the cartilage of the wing of the nose, and expands to the bridge of the nose
5. **LEVATOR LABII SUPERIORIS ALÆQUE NASI** from superior maxillary to cartilage of nose and to the upper lip
6. **ORBICULARIS ORIS**,
 the oval muscle which forms the chief mass of the lips

The following muscles are inserted into the muscular substance of the lips

7. **LEVATOR LABII SUPERIORIS**
 from the superior maxillary and the malar bone to the upper lip
8. **LEVATOR ANGULI ORIS**
 from superior maxillary to corner of mouth
9. **ZYGOMATICUS LABII** } from malar
 MINOR } bone to corner
 10. " MAJOR } of mouth
11. **BUCCINATOR**
 from both superior and inferior maxillary bones to the corner of the mouth
12. **DEPRESSOR ANGULI ORIS**
 from inferior maxillary to corner of mouth
13. **DEPRESSOR LABII INFERIORIS**
 from inferior maxillary to lower lip
14. **LEVATOR MENTI**
 from inferior maxillary to the integument of the chin

MUSCLES OF THE NECK.



The **PLATYSMA MYOIDES** muscle, a thin sheet of muscular fibres arising from the fascia covering upper part of Deltoid and Pectoral muscles. It is inserted above mainly to the skin of the lower part of the face, many of its fibres blending with the muscles about the angle and lower part of the mouth. It is so thin that, except in strong action, or in the aged, it does not obscure the forms of the underlying muscles


The **STERNO-MASTOID** muscle arises by two heads from the Sternum and Clavicle; it is inserted into the mastoid process of the Temporal bone and to the superior curved line of the Occipital bone. The sternal origin is in the form of a rounded tendon; it is separated by an interval from the clavicular origin which is composed of fleshy and aponeurotic fibres

The **STERNO-HYOID** muscle arises from the Clavicle and Sternum and is inserted into the Hyoid bone

The **OMO-HYOID** muscle passes from the upper border of the Scapula to the Hyoid bone. It consists of two fleshy bellies united by a central tendon, which is held in position by fascia attached to the Cartilage of the 1st Rib and to Sternum

The **DIGASTRICUS** muscle consists of two fleshy bellies united by an intermediate, rounded tendon, held in connection with the side of the Hyoid bone by a fibrous loop. The posterior belly arises from the mastoid process of the Temporal bone; the anterior belly arise from the lower jaw-bone


MUSCLES OF THE TRUNK.




The **PECTORALIS MAJOR** muscle arises from the Clavicle, inner third, Sternum, Cartilages of ribs, 2nd to 6th, and from the Aponeurosis of the External oblique muscle
It is inserted into the Humerus, at the outer lip of the bicipital groove

The **DELTOID** muscle arises from the Scapula, spine and acromion process, Clavicle, outer third, and is inserted into the Humerus at the V-shaped deltoid impression

MUSCLES OF THE TRUNK.



FRONT VIEW



In the above diagram the axillary border of the Scapula is moved away from the side of the trunk to show the complete muscle

The **SERRATUS MAGNUS** muscle arises from the upper 8 Ribs and the intervening Costal fascia, and is inserted into the base or vertebral border of the Scapula

The muscle lies between the Scapula and the ribs.

NOTE—


The serrated markings of the superficial part of this muscle lie in a curved line, with convexity downwards, which being continued would pass through the nipple.

The **RECTUS ABDOMINIS** is attached above to the Cartilages of the 5th, 6th, and 7th Ribs, and to the tip of the Sternum.

It is attached below to the Pelvis at the Pubic crest and Pubic symphysis

The tendinous intersections of the Rectus are marked on the diagram a. b. c.

MUSCLES OF THE TRUNK.



The EXTERNAL OBLIQUE muscle is attached above to the lower 8 Ribs by 8 digitations; below, it is attached to the Pelvis at the anterior third of the Iliac crest, and to the Aponeurosis which covers the front of the Abdomen and encloses the Rectus

This Aponeurosis is connected above with the lower border of the Pectoralis major muscle; below, the part of the aponeurosis which extends between the anterior superior iliac spine and the Pubis is called Poupart's ligament. In the median line of the abdomen the interlacing of the aponeuroses of the opposite muscles forms the Linea alba, which extends from the ensiform cartilage or tip of the Sternum to the Pubis


MUSCLES OF THE TRUNK

The TRAPEZIUS muscle



arises from the following parts in the median line of the back of the neck and trunk :—

Occipital bone, protuberance and inner third of superior curved line
Ligamentum nuchæ,
Supra-spinous ligament,
7th Cervical vertebra,
Dorsal vertebrae, spinous processes of all ;

and is inserted into the spine of the Scapula and the outer third of the Clavicle



MUSCLES OF THE TRUNK.



Back view
of
Scapula

Muscles covered by Trapezus
A.—RHOMBOIDEUS MAJOR
B.— " " MINOR } arise

from the spines of the vertebrae, 7th Cervical to 4th or 5th Dorsal, and are inserted into the base of the Scapula.

C.—LEVATOR ANGULI SCAPULÆ arises from the transverse processes of the upper 4 Cervical vertebrae and is inserted into the upper part of the base of the Scapula.

D.)—SPLENIUS arises from the ligamentum nuchae and from the spines of the vertebrae, 7th Cervical and 6 upper Dorsal. It divides into the Splenius Capitis (D), inserted into mastoid process of Temporal bone, and Splenius Colii (D') inserted into transverse processes of 2 or 3 upper Cervical vertebra.

Muscles which are superficial in the triangular interval bounded by the Trapezius, Deltoid, and Latissimus Dorsi.

INFRA-SPINATUS arises from the back of the Scapula and from the strong fascia which covers the muscle. It is inserted into the back of the great tuberosity of the Humerus, uniting its tendon with that of the Teres minor.


TERES MINOR arises from the back of the Scapula at its axillary border and is inserted into the Humerus immediately below the Infra-spinatus tendon.

TERES MAJOR arises from the back of the inferior angle of the Scapula and, passing to the front of the Humerus, is inserted into the inner lip of the bicipital groove.

MUSCLES OF THE TRUNK.

The **LATISSIMUS DORIS** muscle arises from the
 Dorsal vertebrae, spines of lower 5 or 6,
 Lumbar " spines of all,
 Sacral " upper 2 or 3,
 the Supra-spinous ligament,
 the Pelvis posterior third of iliac crest,
 the Ribs, lower 3 or 4.

It is inserted into the Humerus at the bottom of the bicipital groove by a short flat tendon which passes in front of, and partly blends with the tendon of the *Teres major*.



THREE-QUARTER BACK VIEW

The following muscles, which lie beneath the *Latissimus dorsi*, are more or less revealed through its thinner parts:—


ERECTOR SPINÆ,
SERRATUS POSTICUS INFERIOR,
SERRATUS MAGNUS.

The anterior border of the *Latissimus dorsi* gives a distinct surface form, and at its upper part it combines with the *Teres major*, forming the posterior border of the Axilla or armpit.

The **SERRATUS POSTICUS INFERIOR**

(see complete view of muscles of the back) overlies part of the *Erector spinæ* and is covered by the *Latissimus dorsi*. It arises by a thin aponeurosis from the spines of 3 or 4 of the vertebrae, dorsal and lumbar, and divides into 4 flat digitations, which are inserted into 3 or 4 of the lower ribs. The vertical line of its fleshy digitations may influence surface form when the muscle is in strong action.

MUSCLES OF THE TRUNK.



BACK VIEW

References

- a. Aponeurosis of Erector spinae
- A. Sacro-lumbalis, the outer part of the Erector spinae
- B. Longissimus dorsi, the inner part of the Erector spinae
- C. Spinalis dorsi; blends with the longissimus dorsi
- D. Splenius capitis (of the head)
- D'. Splenius colli (of the Neck)
- E. Int-rcostal muscles, filling the spaces between the ribs


The **ERECTOR SPINÆ** arises from the **Sacro-iliac** groove, and from a broad tendon or aponeurosis which covers the lower part of the muscle and stretches across from the spines of the Lumbar vertebrae and sacrum, to the hinder fourth of the Iliac crest

The muscular mass divides into two parts at the level of the lowest rib—an outer part called the **Sacro lumbalis** inserted into the angles of the ribs; and an inner part, called the **Longissimus dorsi**, inserted into the transverse processes of the Dorsal vertebrae and also to the Ribs, between their tubercles and angles

(The **Sacro lumbalis** is also named **ILIO COSTALIS**, being attached to the Ilium below and the Ribs above)

The Erector spinae muscles, with their complicated accessory muscles and prolongations into the neck, fill in the grooves seen in an articulated skeleton, between the spines of the vertebrae and the angles of the ribs on either side. Although covered by the superficial muscles, these fleshy columns have a decided influence on the surface form, more especially in the lumbar region of the back.

MUSCLES OF THE THIGH—FRONT VIEW.



Flexors of the THIGH on the body
 The **ILIACUS**, } from the Pelvis at the iliac crest
 and } and iliac fossa
PSOAS, } from the vertebrae are inserted
 together into the lesser trochanter
 of the Femur

Adductors of the THIGH

The **PECTINEUS**, from the Pelvis, at the ilio-pectineal line, to the back of the Femur
 The **ADDUCTOR LONGUS** from the Pubic portion of the Pelvis to the linea aspera, or rough line, on the back of the Femur
 The **ADDUCTOR MAGNUS** from the Pubis and Ischium it is inserted into the whole length of the linea aspera. The internal portion of the muscle terminates in a tendon attached to the Adductor tubercle on the inner condyle of the Femur

The QUADRICEPS EXTENSOR
 The **RECTUS FEMORIS** arises by two tendons from the Pelvis, at the Anterior inferior iliac spine, and from a groove over the acetabulum

The **VASTUS EXTERNUS** arises from the Femur at the great trochanter and along the outer lip of the linea aspera, or rough line, on the back of the Femur

The **VASTUS INTERNUS** } from the inner lip
 and } of linea aspera and
CRUREUS } from almost the
 (deep seated, being covered } whole length of
 by the Rectus) } front and inner side
 of shaft of Femur

These four muscles are, together, called the **Quadriceps extensor**. Their tendons uniting below, are inserted into the Patella, and are continued by the ligamentum patellae to be ultimately attached to the tubercle of the Tibia

The Patella may be regarded as a sesamoid bone developed in the tendon of the Quadriceps extensor. Note that the fleshy fibres of the Vastus internus descend lower than those of the Vastus externus

MUSCLES OF THE THIGH.—FRONT VIEW.


SARTORIUS and GRACILIS

Diagram of muscles on front of thigh

Mnemonic of the arrangement of muscles on front of thigh, the cross line of the letter N representing the Sartorius



Diagram of front view of lower limb with suggested lines of construction




S. Insertion of Sartorius
G. " " Gracilis

The SARTORIUS muscle arises from the anterior superior iliac spine, and is inserted by an aponeurosis into the upper part of the inner surface of the Tibia

The GRACILIS arises from the margin of the Pubic symphysis, and descending ramus of the Pubis. It tapers down to a long tendon which passes close behind the Sartorius, to be inserted into the Tibia

NOTE.—The blending of the line of the Sartorius muscle with the subcutaneous surface of the Tibia forms an important running line in drawing the front view of the leg

MUSCLES OF THE BUTTOCK AND HIP.




The **GLUTEUS MAXIMUS** muscle arises from the posterior fourth of the iliac crest, from the Sacrum and Coccyx, from the aponeurosis of the Erector spinæ muscle, and from the great Sacro-sciatic ligament. The deeper fibres of the lower half of the muscle are inserted into the linea aspera on the back of the Femur; the fibres of the upper half and the superficial fibres of the lower portion, terminate in a strong tendinous lamina which passes across the great trochanter and is inserted into the fascia lata of the thigh

The **GLUTEUS MEDIUS** muscle (partly covered by the G. maximus; and its superficial portion lying between G. maximus and the Tensor fasciæ femoris, covered by a strong fascia) arises from the iliac crest and part of the outer surface of the Ilium. It converges to a strong flattened tendon, which is inserted into the great trochanter of the Femur

The **TENSOR VAGINÆ** or **FASCIÆ FEMORIS** arises from the iliac crest close to the antr. supr. iliac spine. It is inserted into the fascia lata

MUSCLES OF THE THIGH—BACK VIEW.




The HAMSTRING muscles, flexors of the leg

The **BICEPS** (the outer hamstring) arises by two heads; the long head from the back of the tuberosity of the Ischium by a tendon common to it and the Semitendinosus; the short head from part of the linea aspera on the back of the Femur. It is inserted into the head of the Fibula with an expansion of the tendon to the Tibia.

The **SEMITENDINOSUS** (together with the semimembranosus forming the inner hamstring), arises from the tuberosity of the Ischium. It tapers below to a very long tendon inserted into the upper part of the inner surface of the Tibia.

The **SEMIMEMBRANOSUS** (so named from the membranous expansion on its anterior and posterior surfaces) arises from the tuberosity of the Ischium and is inserted into the back of the inner tuberosity of the Tibia.

Front View of the bones at the knee




St. Insertion of
Semitendinosus
G Insertion of Gracilis
S " " Sartorius

MUSCLES OF THE LEG.—BACK VIEW.

Muscles of the calf of the leg

The **GASTROCNEMIUS** muscle arises by two heads from the Femur immediately above the condyles, and ends below in a broad tendon which joins with that of the Soleus to form the Tendo-Achillis. Each tendon of origin spreads out into an aponeurosis from which some of the muscular fibres arise. The two heads meet in the median line of the calf

The **SOLEUS**, a broad flat muscle, shaped like a sole-fish, arises from the back of the Tibia and Fibula. The fleshy fibres are short and pass backward to an aponeurosis which joins below with the tendon of the Gastrocnemius, forming the Tendo-Achillis, which is inserted into the Os calcis or heel-bone




Deep muscles of the back of the leg. These muscles are superficial only at the lower part of the inner border of the leg. They arise from the back of the Tibia and Fibula and from the interosseous membrane, which extends between these two bones and separates the muscles of the front from those on the back of the leg.


The **FLEXOR LONGUS DIGITORUM** terminates below in a tendon which descends behind the inner malleolus along with the T. posticus, and crossing superficially to the tendon of the Flexor longus pollicis, passes into the sole of the foot and divides into four tendons for the outer toes.

The **TIBIALIS POSTICUS** is inserted into the Scaphoid bone, with prolongations to most of the tarsal and metatarsal bones

The **FLEXOR LONGUS POLLICIS** or **HALLUCIS**, almost entirely hidden, is inserted into the base of the last phalanx of the great toe



MUSCLES OF THE LEG—FRONT VIEW.



OUTER VIEW

The **TIBIALIS ANTICUS** arises from the Tibia at its outer tuberosity and outer surface of the shaft, and from the interosseous membrane. Its tendon, after passing through the innermost compartment of the anterior annular ligament, is inserted into the inner cuneiform bone and the base of the metatarsal bone of the great toe

The **EXTENSOR PROPRIUS POLLICIS** or **HALLUCIS** (of the great toe) arises from the front of the Fibula and the interosseous membrane. Its tendon, passing through a compartment of the annular ligament, is inserted into the base of the last phalanx of the great toe



THE **EXTENSOR LONGUS DIGITORUM** arises from the outer tuberosity of the Tibia and the upper three-fourths of the shaft of the Fibula, and from the interosseous membrane. Its tendon passing through the annular ligament divides into four slips which are inserted into the four outer toes at their 2nd and 3rd phalanges

The **PERONEUS TERTIUS** is a part of the Ex. longus digitorum. It arises from the lower fourth of the shaft of the Fibula. The tendon, after passing through the same compartment of the annular ligament as the Ex. longus, is inserted into the base of the metatarsal bone of the little toe

The **PERONEUS LONGUS** arises from the head and upper two-thirds of outer surface of the shaft of the Fibula. It terminates in a long tendon, which, passing behind and beneath the outer malleolus in a groove common to it and the Peroneus brevis, is directed forwards and downwards to the outer border of the foot and enters a groove on the under surface of the Cuboid bone; it then passes deeply across the sole of the foot to be inserted into the inner cuneiform bone and the base of the metatarsal bone of the great toe.



The **PERONEUS BREVIS** lies beneath the Peroneus longus and arises from the lower two-thirds of the outer surface of the shaft of the Fibula. Its tendon passes behind the outer malleolus along with the Peroneus longus, and separating from the latter at that point, turns forward along the outer side of the foot, to be inserted into the tuberosity of the 5th metatarsal bone

OUTER VIEW OF THE FOOT.




The **ABDUCTOR MINIMI DIGITI** arises from the os calcis and is inserted into the 1st phalanx of the little toe. It is slightly attached in its course to the base of the 5th metatarsal bone. The **EXTENSOR BREVIS DIGITORUM** arises from the os calcis and sends tendons to the four inner toes. This muscle causes an important surface form in front of the outer ankle.

INNER VIEW OF THE FOOT.



The **ABDUCTOR POLLICIS** arises from the os calcis and is inserted into the internal sesamoid bone and the 1st phalanx of the great toe

MUSCLES OF THE ARM.—FRONT VIEW




The **BICEPS** muscle is divided above into two portions or heads. The short head arises from the coracoid process of the Scapula along with the Coraco-brachialis. The long head arises by a long and rounded tendon, from the upper margin of the glenoid cavity of the Scapula, the socket of the shoulder-joint. This tendon passes over the head of the Humerus and lies in the bicipital groove, in which it is held by an expansion of the Pectoralis major tendon. The two portions of the muscle join about the middle of the arm, and the muscular mass terminates above the elbow in a flattened tendon inserted into the back part of the tuberosity of the Radius. Opposite the bend of the elbow the tendon gives off from its inner side, a broad aponeurosis, the Bicipital fascia, which is continuous with the deep fascia or sheath of the muscles of the forearm.

The **CORACO-BRACHIALIS** muscle arises from the coracoid process of the Scapula along with the short head of the Biceps. It is inserted into the Humerus at the middle of the inner surface of the shaft, between the origin of the Triceps and Brachialis Anticus.

The **BRACHIALIS ANTICUS** is a broad muscle which covers the lower half of the front of the Humerus and the front of the elbow joint. It arises from the lower half of the front of the shaft of the Humerus, commencing above at the insertion of the Deltoid, which it embraces by two angular processes. Its fibres converge below to a thick tendon which is inserted into the coronoid process of the Ulna.

MUSCLES OF THE ARM — BACK VIEW.



The TRICEPS muscle, situated on the back of the arm, extends the entire length of the back of the Humerus. It is divided above into three parts, named :

1. The middle, or long head,
2. The external head,
3. The internal, or short head.

The middle or long head arises from a depression immediately below the glenoid cavity of the Scapula. The external head arises from the upper and outer part of the back of the Humerus. The internal head arises from the back of the Humerus and from its inner border, commencing above narrow and pointed and extending below near to the trochlear surface. The *common tendon* of the Triceps receives the muscular fibres from the three heads; it commences about the middle of the back of the muscle and causes a very important flattened surface form. It is inserted below into the olecranon process of the Ulna, an expansion of the tendon being continued downwards on the outer side, passing over the Anconeus muscle, to blend with the deep fascia of the forearm. The long head of the Triceps passes between the Teres minor and Teres major muscles.

The ANCONEUS muscle is a small triangular muscle, placed behind and below the elbow joint. It appears to be a continuation of the outer portion of the Triceps. It arises from the back of the outer condyle of the Humerus, and is inserted into the side of the olecranon process and the upper fourth of the back of the shaft of the Ulna.




Diagram of the surface forms on the back of the arm

MUSCLES OF FRONT AND INNER SIDE OF FOREARM. THE PRONATOR AND FLEXOR GROUP.




Diagram of the four superficial muscles of the front of the forearm



The Flexors of the fingers lying deep in the forearm and covered by the flexors of the wrist, except in the interval between the tendons of the Palmaris longus and the Flexor carpi ulnaris

The **PRONATOR and FLEXOR** group. These muscles, which are situated on the front and inner side of the forearm, arise from the internal condyle of the Humerus by a common tendon


The **PRONATOR TERES** muscle passes obliquely across the forearm to its insertion at the middle of the outer surface of the shaft of the Radius

The **FLEXOR CARPI RADIALIS** muscle terminates below in a tendon which forms the lower two-thirds of its length. It is inserted into the base of the metacarpal bone of the index finger, passing through the annular ligament and through a groove (which is enclosed by a fibrous sheath) in the Trapezium bone of the wrist

The **PALMARIS LONGUS** muscle terminates in a slender flattened tendon which passes over the annular ligament and ends in the Palmar fascia

The **FLEXOR CARPI ULNARIS** muscle terminates in a tendon which occupies the anterior part of the lower half of the muscle, and is inserted into the pisiform bone, being further prolonged by ligaments to the 5th metacarpal bone and the ulnar bone. Besides its attachment above to the inner condyle of the Humerus, this muscle also arises from the olecranon process and from the upper two-thirds of the posterior border of the Ulna

MUSCLES OF BACK AND OUTER SIDE OF FOREARM. THE SUPINATOR AND EXTENSOR GROUP.



The SUPINATOR and EXTENSOR group

These muscles lie on the outer side and back of the forearm. It is important to note that they form a mass which commences on a higher level than that of the Pronator and Flexor group. The apex of this muscular mass arises from the Humerus, some distance above the elbow joint, emerging from between the Triceps and Brachialis anticus. The Supinator longus and the Extensor carpi radialis longior arise from the ridge above the outer condyle of the Humerus, and the rest of the group take origin from the outer condyle. The Extensors of the thumb form a separate group arising from the Radius and Ulna.

The SUPINATOR LONGUS, fleshy in its upper two thirds, terminates in a flat tendon inserted in styloid process of Radius.


The EXTENSOR CARPI RADIALIS LONGIOR terminates at the upper third of the forearm in a flat tendon which passes beneath the extensors of the thumb, and through a groove behind styloid process of Radius, to be inserted into base of metacarpal bone of index finger.

The EXTENSOR CARPI RADIALIS BREVIOR tendon begins about the middle of the forearm and accompanies that of the preceding muscle. It is inserted into the base of the metacarpal bone of the middle finger.


The EXTENSOR COMMUNIS DIGITORUM divides just below the middle of the forearm into three tendons, which pass through the annular ligament and then diverge, the innermost one dividing into two. These tendons are inserted into the second and third phalanges of the fingers.

The EXTENSOR MINIMI DIGITI a slender muscle connected with the preceding. Its tendon, after passing through the annular ligament divides into two, which, again uniting are inserted into the 2nd and 3rd phalanges of the little finger.

The EXTENSOR CARPI ULNARIS. This muscle, besides the common ridge from the Humerus, also arises by an aponeurosis from the posterior border of the Ulna. Its tendon runs through a groove behind the styloid process of the Ulna, passes through the annular ligament, and is inserted into the base of the metacarpal bone of the little finger.



BONES OF THE HAND.—PALMAR VIEW



MUSCLES OF THE PALM

- Muscles of the Thumb forming the Thenar eminence or ball of the thumb**
1. OPPONENS POLLICIS
 2. FLEXOR BREVIS POLLICIS
 3. ABDUCTOR POLLICIS
 4. ADDUCTOR } transversus and
 } obliquus

- Muscles of the Little Finger, forming the Hypothenar eminence or ball of little finger**
1. ABDUCTOR MINIMI DIGITI
 2. FLEXOR BREVIS " "
 3. OPPONENS " "

The ABDUCTOR MINIMI DIGITI ..
The FLEXOR BREVIS ..
The OPPONENS ..

Origin


From the Pisiform bone
from unciform process of
unciform bone and from
the annular ligament

Insertion

1st phalanx of little finger
do. do.
Metacarpal bone

In the middle of the Palm are the Lumbricales, four small muscles, accessories to the deep Flexor of the fingers; and in the intervals between the metacarpal bones, are the Interossei muscles, 4 dorsal and 3 palmar

MUSCLES OF THE HAND.—PALMAR VIEW.



The THUMB is provided with the following muscles :—
 3 Extensors situated on the back of the forearm and hand ;
 4 Flexors, 1 Abductor and 2 Adductors, all muscles of the palm, except the Flexor Longus, which has its fleshy part deep in the front of the forearm

EXTENSORS

EX. OSSIS METACARPI POLLICIS inserted into the
 " PRIMI INTERNODII POLLICIS " " "
 " SECUNDI INTERNODII POLLICIS " " "

Metacarpal bone
 1st Phalanx
 terminal Phalanx

FLEXORS

OPONENS POLLICIS arises from the Trapezium and annular ligament and is inserted into the

Metacarpal bone

FLEXOR BREVIS POLLICIS consists of two portions ; the outer arises from the Trapezium and annular ligament and is inserted into the

1st Phalanx, outer side

The inner and deeper portion arises from the metacarpal bone of the thumb and is inserted along with the Adductor obliquus into the

1st Phalanx, inner side

A sesamoid bone is developed in each of the two tendons of insertion

FLEXOR LONGUS POLLICIS, a deep muscle of the forearm having an extensive origin from the Radius ; it is inserted into the

terminal Phalanx

ABDUCTOR and ADDUCTORS



ABDUCTOR POLLICIS arises from Trapezium and annular ligament and is inserted into the

1st Phalanx



ADDUCTOR POLLICIS } obliquus, from the Os magnum, etc.,
 transversus, from the metacarpal bone of the middle finger

inserted into
 1st Phalanx


BONES OF THE HAND.—BACK VIEW



BACK OF THE HAND—MUSCLES.




SURFACE FORMS OF THE LIMBS.




INNER VIEW OF BENT KNEE

OUTER VIEW OF BENT KNEE




Front view of knee

Back view of knee



INNER VIEW OF FOOT

OUTER VIEW OF FOOT




BONES OF THE UPPER AND LOWER LIMBS.

Bones of the
UPPER LIMB,
front view, with the forearm
in the position of
PRONATION

NOTE.—The illustrations on
this page, and also those of
the skeleton of the trunk in
three-quarter front and back
views, are from photographs
of artificially articulated
bones

Bones of the
LOWER LIMB
in bent position
Outer view





Bones of the
UPPER LIMB,
front view, with the
forearm in the posi-
tion of
SUPINATION

NOTE.—The bones of
the upper limb are
here shown on a
larger scale than those
of the lower limb


Bones of the
LOWER LIMB
in bent position
Front view



DIAGRAMMATIC NOTES OF VARIOUS BONES




SURFACE FORMS OF THE TRUNK.



**THREE-QUARTER
FRONT AND BACK
VIEWS OF THE
MALE TORSO OR
TRUNK.**




**THREE-QUARTER
FRONT AND BACK
VIEWS OF THE
FEMALE TORSO.**



**DIAGRAMS SHOWING
THE PELVIC BONES AND
THE THIGH BONES IN
BACK VIEW IN THE
MALE AND FEMALE.**




**DIAGRAMS OF
SIDE VIEW OF
PELVIC BONES
SHOWING INCLINATION
IN MALE AND FEMALE.**



**MALE
PELVIS.**


**FEMALE
PELVIS.**




**BACK VIEW
IN THE MALE.**

**BACK VIEW
IN THE FEMALE.**


THREE-QUARTER FRONT VIEW OF SKELETON.




THREE-QUARTER BACK VIEW OF SKELETON.



DIAGRAMS WITH NOTES ON PROPORTION.



Relation of Head to height of figure.
 At Birth. 5 years. Adult.



In the adult male figure, in the upright position, the distance from the top of the head to the pit of the neck measures about $5\frac{1}{2}$ times into the height; this distance is about equal to the greatest width at the hips, or middle of the height of the figure. The greatest width at the shoulders, at the fullest point of the deltoid muscle, is about equal to 2 heads, or more than one quarter of the height.

In the female figure, the distance from the top of the head to the waist is about $\frac{1}{3}$ of the height. The width at the shoulders is slightly less proportionately than in the male figure, but the width at the hips is proportionately, or even absolutely, greater in the female, and is at a lower level, being a little below the great trochanters. The width at the hips measures about $4\frac{1}{2}$ times into the height, and is about equal to the distance from the pit of the neck to the umbilicus.



FA6478.5

Anatomical diagrams for the use of
Fine Arts Library **AZ70632**



3 2044 034 097 758

